

**U-CyTech B.V.**  
Yalelaan 48  
3584 CM Utrecht  
The Netherlands  
+31 85 073 1460  
info@ucytech.com  
www.ucytech.com

<b>Data sheet</b>	<b>Monoclonal antibody to human interleukin (IL) 12 / IL-23 p40 (IL-12/IL-23 p40)</b>
<b>Cat. No</b>	<b>CT257</b>
Clone	B-P24
Isotype	Mouse IgG <sub>1</sub> , κ
Immunogen	Recombinant human IL-12
Production	<i>In vitro</i> using serum free medium
Purification	Ion exchange chromatography
Packaging	Lyophilized and vacuum-packed
Contents	0.5 mg/vial
Buffer	Prior to lyophilization: 0.5 ml PBS + 125 mM trehalose.
Specificity	Binds both natural and recombinant human IL-12 (p35/p40) and IL-23 (p19/p40) heterodimers via the p40 subunit <i>in vitro</i> . Blocks IL-12 induced proliferation of PHA and IL-2 activated lymphocytes.
Sterility	Membrane filtered (0.2 μm)
Reconstitution	Dissolve the contents of the vial by injection of 0.5 ml sterile distilled water.
Stability	Lyophilized product is stable for at least one year at 2 to 8°C (expiry date is printed on the vial). After reconstitution, the contents can be safely stored at 2 to 8°C for one month or for one year at -32 to -18°C. Add 0.02% sodium azide to prevent bacterial growth.
Quantitation	Antibody concentration was determined by absorbance, taking A <sub>280</sub> =1.4 for a 1 mg/ml solution.
Application	ELISA and ELISPOT assays Intracellular staining <i>In vitro</i> neutralization

References can be found in the Reference Database at [www.ucytech.com/references](http://www.ucytech.com/references).

